Monthly Update Technical Study in the Relationships of Solar Flux, Water, Carbon Dioxide To Global Weather Patterns

David Pristash
Independent Researcher

October, 2022 Data

The purpose for this paper

This paper is not meant to be a peer-reviewed work; but it is meant to give a foundation for a more serious study of the subject matter presented here which is of determining the basis of developing a global temperature. There are three areas of interest.

- 1) The amount of thermal energy that reaches the planet from the sun.
- 2) The amount of thermal energy that is initially absorbed by the planet.
- 3) The process on the planet that 'temporarily' holds thermal energy on the planet.

In this paper, I will give a frame work for determining all three aspects.

Part One, the blackbody temperature of the planet
Part Two, the planetary greenhouse effect
Part Three, the probable range of temperatures on the planet

Appendix

NASA Table Land Ocean Temperature Index (LOTI) April 2008

NASA Table Land Ocean Temperature Index (LOTI) current to the date of this paper

Part One, the Blackbody Temperature of the Planet Earth

Determining the 'exact' blackbody temperature of the planet is the first step in determining what the "greenhouse' effect is; for without that value all else is either speculation or based on an unreliable value. This leads us to a quandary since the planet is a globe spinning around a titled axis of rotation and with an elliptical orbit around the sun **Figure 1** which is the source of virtually all the energy that heats the planet. Clearly, with these facts there cannot be one temperature for the planet and so, in theory, an average could be calculated but it will also be very misleading and lead to false conclusions; especially as it hides very large energy flows on the planet.

Traditional calculations of the planets black body temperature ignore the variables which then lead one to assume a steady state situation verses the real dynamic situation that actually drives climate. To justify this assumption a general statement that the variances are too small to have any meaningful effect are promoted. In some cases, maybe with fewer variables, this might be true but in this case, I think not.

These are the main variables, constants, and forces:

- 1. The sun has a primary and secondary cycle the primary is Magnetic of about 22 years (Pi times 7) which changes the polarity of the suns magnetic field which therefore gives a variation in the suns solar wind which is the more important.
- 2. The secondary cycle is the number of sun spots which is half the magnetic at about 11 years and that gives a small variation in the suns output of about 1%
- 3. The planet has an elliptical Orbit that varies by 3.34% or 4,999,849 miles, at this time.
- 4. The axial tilt of the planet is 23.4 degrees which causes winter and summer to alternate between Aphelion and Perihelion about every 10,000 years
- 5. The planet is a sphere so only one side faces the sun at any given moment
- 6. The sun's energy reaches the planet on a line drawn from the center of the sun to the center of the planet which only intersects the equator twice a year.
- 7. Actually, the line from the sun to the earth is to the barycenter (center of mass) of the earth and the moon system. Which changes the distance to the sun to the earth's surface by +/-2,858 miles per lunar month; however, this complication is ignored in the study.
- 8. The energy from the sun is concentrated around this line, a hot spot.
- 9. The planet is a sphere so the suns radiation drops off in all directions from this line by a Cosine factor to zero at the edge 90 degrees from the center line
- 10. The spin and tilt of the planet means that the center line, in effect, moves up 23.4 degrees from the equator and then down 23.4 degrees from the equator during the course of one orbit
- 11. That movement means the distribution of the energy in the hot spot also moves
- 12. The distribution of land and ocean are not uniform on the planet and therefore the absorption of the solar flux is very different at points the hot spot travels over.
- 13. The albedo of the planet is a variable not a constant mainly as a factor of the amount and kind of clouds.
- 14. Energy from the core adds a small amount of energy, but not enough to matter.

- 15. Tidal forces from the sun and the moon also add some energy, again to small to matter
- 16. Energy is carried North and South from the hot spot, centered on the line described in item 6, by the atmosphere and the ocean
- 17. The Coriolis Effect along with tidal forces drive thermal transfer north and south at an angle and these are the main contributors to the climate





Figure 1, the Earth's orbit



There are three sources of energy that determine the climate on the earth: the radiation from the sun which is said to be 1366 Wm² The actual value based on the orbital range is from 1414.4 Wm² in January to 1323.0 Wm² in July see **Chart 13** and there is also an eleven year sun spot cycle with a range of 1.37 Wm². The hot core of the planet adds ~0.087 W/m² and the gravitational effects of the moon and the sun (tides) adds another ~.00738 Wm². Of these three, the sun's radiation is by far the most important but considering all three the range during an eleven year solar cycle is from a high of ~1415.3 Wm² to a low of ~1322.4 Wm² so a more accurate mean would be 1368.34 Wm².

The energy emitted by the planet must equal the energy absorbed by the planet and we can calculate this using the Stefan-Boltzmann Law. Which is the energy flux emitted by a blackbody is related to the fourth power of the body's absolute temperature. In the following example, the tidal and core temperatures are added after the albedo adjustment since they are not reduced by the albedo.

Earth's blackbody temperature

 $= \sigma T^4$

Ε

Earth's surface temperature (when report first written)

```
T_{bb} = 252.23° K (-20.92° C) low T_{s} = ~287.75° K (14.6° C) today T_{bb} = 254.36° K (-18.79° C) mean T_{bb} = 256.54° K (-16.51° C) high
```

The difference between the blackbody and the current temperatures is what we call the 'greenhouse' effect that averages 33.36° Celsius (C), today, although the range is from 35.52° C to 31.11° C from variations in the 11 year solar cycle. This documented variation means that the stated Blackbody radiation as shown here will give a 4.41° C variation or let's say 14.0° C plus or minus 2.2° C because of the Stefan-Boltzmann Law which has a 4th power amplification. This will result in a slow 11 year cycling fluctuation of energy in the tropics where the bulk of the energy comes in that is not inconsequential.

If we add clouds to the picture it get even more complex as they have a significant effect on the planets albedo as we know from two major volcanoes' both in Indonesia; one in 1815 Tambora and the other in 1883 Krakatoa both of which threw enough particles into the atmosphere to significantly lower the temperature of the planet. Although dust is not a cloud the point is that if the albedo of the

planet is changed it does have a major effect on global temperatures. The lack of thermometers in 1815 means we really don't know what the effect was other than 1816 is known as the year without a summer. The other eruption in 1883 is well documented and is estimated to have dropped world temperatures by 1.20° C which would be equivalent to about a 4.2% reduction in the global albedo. The importance of clouds can be seen in the following Chart Figure 3. A reasonably estimate of the total effect of clouds on the global albedo would be about 50% if nothing else changed or a reduction in Albedo from 30% to 15%. Just for reference the Albedo of the moon is 13.6% which if the earth had no clouds or water but still had an atmosphere the black body temperature of the planet would be 268.71° K or -4.4° Celsius.




Figure 3, Albedo of various surfaces

Just for sake of argument if we varied the cloud levels by +/- 10% we find that at low solar flux and high clouds the Blackbody temperature would be 249.46° K and with high solar flux and low clouds the Blackbody temperature would be 259.32° K a range of 9.86° C. The reason this is so important is that properly modeling cloud levels is the area with the most uncertainly in the present climate models as clouds form at much lower mesh resolutions (an aspect of the process used) then the present models can deal with even if the formation could be properly modeled.

Despite this variation in incoming solar flux the planet's temperatures has been very stable as shown in **Figure 4** so we know there are no positive or negative feedback process of any consequence on the planet that would create a runaway temperature scenario. Other factors are also important in doing climate work such as 52.3% of the solar energy is concentrated within 45.0 degrees of the hot

spot and 77.6% within 60 degrees of the hot spot. And the heat from the core and probably the tides is concentrated where the crust is the thinnest under the oceans and this concentration of energy core heat and tides) combined with Coriolis forces is probably what drives the ocean currents. In my opinion these other important factors are not being considered properly in the climate models, and that results in climate models that don't work properly e.g. the inability to explain why there has been a pause in the warming calculated by NASA and NOAA over that past ten years despite a continuing increase in the level of CO₂ in the atmosphere.

We also know from geological studies **Figure 4** that the planets temperature has been relatively stable over the past 600 million years with a mean of about 17° C or 290° Kelvin (K) and with a range of plus or minus 5° K or C based on the information in **Figure 4**. During the past 250 million years CO_2 concentrations have ranged from a low of ~280 ppm (a historic low) in 1800's to the present low of 410 ppm to a high of over 2,000 ppm probably averaging around 1,500 ppm. There was only one other period in the past 600 million years with CO_2 this low. Going back, further CO_2 was estimated to be as high as 7,000 ppm, but we will ignore that for now.

This means that whatever the processes are that relate to determining the thermal balance of the planet they must work within this range of ~12° C to be valid. Although **Figure 4** shows a range of 10° C it would be prudent to spend resources to determine these values with as great accuracy as possible. We'll suggest a mean of 17° C with a range from 10° to 24° C as being more reasonable in this work. Also, we are now in one of only three cold periods which are very rare in the past 600 million years and if we count that partial dip 150 million years ago that means that there is probably a 150 million year cycle there; maybe one of those first determined my Milutin Milankovic.




Figure 4, Geological temperatures, and Carbon Dioxide

Additional discussion as to the so called "greenhouse" effect must start with the important correction that this process is not a true greenhouse effect, since it is not the same process that occurs in a greenhouse used to grow food, not even remotely close to it.

The actual process that occurs is based on the structure of the atoms involved and how they interact with the various frequencies of visible and infrared radiation that are in play on the planet after arriving here from the sun. However at this point in time there is no way to correct for the misuse of the words so we are stuck with it and all the complications that therefore arise in trying to properly discuss the issue with lay people and even some with technical knowledge.

The greenhouse effect occurs within the earth's atmosphere and the main constitutes of wet air, by volume ppmv (parts per million by volume) are listed in the following table. Water vapor is 0.25% over the full atmosphere but locally it can be 0.001% to 5% depending on local conditions. Water and CO₂ are mostly near the surface not in the upper atmosphere so the bulk of the greenhouse effect must be close to the surface. This table is slightly different than most as it shows water.

Gas	Volume	Percentage
Nitrogen (N ₂)	780,840 ppmv	78.8842%
Oxygen (O ₂)	209,460 ppmv	20.8924%
Argon (Ar)	9,340 ppmv	0.9316%
Water vapor (H ₂ O)	2,500 ppmv	0.2494%
Carbon dioxide (CO ₂)	400 ppmv	0.0399%
Neon (Ne)	18.18 ppmv	0.001813%
Helium (He)	5.24 ppmv	0.000523%
Methane (CH ₄)	1.79 ppmv	0.000179%

There are only two of these gases that are relevant to determining how that 33° C (today) happens. That is not to say the others do not contribute but that at the present concentrations of Water H₂O and Carbon Dioxide CO₂ they are the main determinants. And since we know the range of temperatures that have existed geologically then we have set the range which these two gases must interact in, meaning that any set of equations, models, or theories that predict values outside this range must be suspect based on geological evidence.

Also, it must be kept in mind that the solar flux falls on a spot centered on a line drawn from the center of the earth to the center of the sun and because of the 23.4° axial tilt of the planet, this "Hot" spot moves up and down as the planet moves through its orbit. Because of the shape of the planet, the intensity falls off quickly as we move north, south, east, and west according to a cosine factor so the heat energy is mostly over oceans near the equator where the atmosphere is the densest.

The first image below **Figure 5** shows a recent distribution of water across the planet, it is clearly concentrated over the oceans close to the equator, and that results in the heat imbalance and therefore movement north and south as shown in the second image **Figure 6**.






Figure 5, water vapor concentrated near the equator





In summary we now know that the Blackbody temperature of the planet is a variable.

```
T_{bbl} = 252.23° K (-20.92° C) low at Aphelion T_{bbm} = 254.36° K (-18.79° C) and the yearly mean T_{bbh} = 256.54° K (-16.51° C) high at Perihelion
```

Therefore the 'greenhouse effect, with clouds as a constant, must be a variable.

$$T_s = \sim 287.75^{\circ} \text{ K } (14.6^{\circ} \text{ C}) \text{ today}$$

$$Gh_l = T_{bbl+} T_s = 35.52^{\circ} \text{ C}$$

$$Gh_m = T_{bbm+} T_s = 32.39^{\circ} \text{ C}$$

$$Gh_h = T_{bbh+} T_s = 31.11^{\circ} \text{ C}$$

Considering there would probably be fewer clouds during cool period and more clouds during warm period the following would be more like the true effect considering both.

```
T_{bblc}=252.98^{o} K (-20.17° C) low at Aphelion T_{bbmc}=254.36^{o} K (-18.79° C) and the yearly mean T_{bbhc}=255.83^{o} K (-17.32° C) high at Perihelion
```

Therefore the 'greenhouse effect with clouds included must also be a variable. In this case we assume fewer clouds in cooler periods and more clouds in warmer periods of 2.5% which reduces the range and acts as a negative feedback on the process.

$$T_s = \sim 287.75^{\circ} \text{ K } (14.6^{\circ} \text{ C}) \text{ today}$$

$$Gh_{lc} = T_{bblc} + T_s = 34.77^{\circ} \text{ C}$$


$$Gh_{mc} = T_{bbmc} + T_s = 32.39^{\circ} \text{ C}$$

$$Gh_{hc} = T_{bbhc} + T_s = 31.92^{\circ} \text{ C}$$

Chart 16 on the next page shows the true black body temperature of the planet over the period of one year assuming an atmosphere with no water or no CO₂. The blue plot is the actual and the yellow plot is the generally accepted average. The blue curve is plotted from the distance to the sun of the planet and accepted output of the sun in Wm² of the sun. Because of the Stefan-Boltzmann Law the small change in solar radiation, reaching the planet from the sun is magnified by the 4th power such that it really does make a difference as is clearly shown in **Chart 16**.


A swing of 4 degrees Celsius cannot be ignored when developing a climate model, especially when we are talking climate changes of a few degrees from CO₂ that are going to melt the planet. I cannot understand what has happened to science today it's like a belief in some bizarre pagan god demanding a sacrifice of a virgin girl every day to prevent the rest of us from being consumed by the environmental god Al Gore.

Chart 16, Black Body Temperature of planet -18.76 degrees Celsius



The next Chart, **Chrat 17** adds to **Chart 16** he NASA global temperatures from three years 2007, 2012 and 2017. The Black Body temperature and the NASA plots are not shown together because it will show a problem. If the planets black body temperature is dropping in the summer, as it must by orbital mechanics, then how can the NASA global temperatures remain constant? The issue that this Chart shows results from NASA using an average value for solar flux rather than the true value.

Chart 17, Black Body Temperature Verses Monthly LOTI Values



One more Chart, **Chart 18** shows the difference between the black Body temperature shown in **Chart 16** and **Chart 17** subtracted from the NASA temperatures. This seems to show that there is some mysterious energy that enters the planet atmosphere to add almost 5 degrees Celsius to the planet during the Aphelion phase of the planets orbit. Clearly, something is wrong here and I am reasonably sure that it is the homogenization process that NASA used to make up the global temperature.





Chart 18, Black Body Temperature of planet

The range in temperature just from orbital changes is 4.41° C but including clouds that range is reduced to 2.85° C however in either case it is significantly more than the warming that the IPCC claims has happened looking at only Carbon Dioxide as the main factor. These are hard numbers based on the solar flux which is known and the orbital parameters of the Earth that are also known. The large temperature variances come from the Stefan-Boltzmann Law; which is the energy flux emitted by a blackbody is related to the fourth power of the body's absolute temperature (meaning we must use kelvin). The fourth power in the equation magnifies the small variation in solar flux significantly.

With the understanding that we have now o the importance of Black Body temperature we can add two key factors that will determine the range of possible global temperatures based on a mathematical development of the sensitivity values of H_2O and CO_2 . The sensitive of H_2O and CO_2 determines the delay factor of the thermal energy contained in the atmosphere, the oceans, and the land which must leave the planet to put things in balance which is required to happen. Also the warmer the planet gets the faster the heat will leave. These basic principles will allow us to develop a series of curves that represent the various possible temperature ranges of the planet in the last section of this paper.

On the this page, **Figure 8** shows that complexity of the atmospheric energy flows that the global climate models are trying to duplicate in software. Considering that there are no fixed numbers or values and this is a very dynamic situation in which one of the key determinate of "climate" cloud formation has not been modeled it hard for me to see how there is any chance of the models being anything other but a science/programing toy.



© 2005 Brooks/Cole - Thomson

Figure 7 Basic atmospheric flows

Part Two, The Planetary Greenhouse Effect

Now that we have a better understanding of the blackbody temperature of the plant, we can discuss the process that brings us to the global temperature range that we live in. Keep in mind that the temperature of the earth would be the same as that found on the moon if there was not some way to make it warmer. The moon is, after all, the same distance from the sun that are our planet is.

So what is the average temperature of the moon? Using the methods we used to calculate the black body temperature of the earth gives us approximately 236° Kelvin or -37° Celsius which is 18° Celsius colder what we calculated for the earth. So the first thing we learn is that there must be two black body temperatures, so to speak. Considering there is an atmosphere but no life would give us -18.2° Celsius of a stored thermal energy buffer. And then we need another 32.8° Celsius to bring us to the 1950-80 Base NASA temperature 14° Celsius. This makes the earth approximately 51° Celsius warmer than the moon

Therefore, the real thermal buffer is somewhere around 50 ^O Celsius plus or minus a degree or two based on NASA data and that is the amount that needs to be explained by any theories of global climate which is not nor has ever has been a constant. Blaming humans for what is obviously a natural variable is foolish at best and criminal if used for political ends.

From part one, we know that the amount of thermal radiation that reaches the planet has enough variance to it that it needs to be accounted for in any valid theories or calculations. So with that out of the way we'll now look at the so called greenhouse effect which is approximately 33° Celsius. But this brings us to another issue. The process that allows the suns thermal energy to be held in a buffer and warm the planet is modeled as a log function which means that as the variables (H₂O & CO₂) increases the effect of the variable diminishes such that at some point there is no more effect.

Back in July 1979, the US National Academy of Sciences was given the task of determining what that log function looked like and they came up with an Ad Hoc Study Group which issued a report by the end of that year. It was thereafter called the Charney Report as Jule G. Charney was the Chairman of the Ad Hock study group. The key result was that the increasing effects from the doubling of CO_2 were estimated to be from 1.5° C to 4.5° C, or 3.0° C +/- 1.5° C. That ended up being the values that were used to build all the Climate Models used by the IPCC since it did apparently explain the observed changes in global temperatures, at that time.

There were three oversights made at that time.

- 1) There are no naturally occurring climate changes (not true based on geological evidence)
- 2) There was only one peer-reviewed paper on the subject (way too few to define the issue)
- 3) The effect in question could not be a log function (the equation has to be logistic function)

If the first oversight had not been made Issue, two and three would not have occurred because the second oversight produced equations that gave too large a value to the doubling. Years later additional scientific work would indicate that the 3.0° C +/- 1.5° C. was more likely in the range 1.5° C +/- .75 C. That work was ignored as the observed temperature changes could not be reconciled with

the lower values and therefor CO₂ would not be a global problem. Since the geological temperatures showed the Charney report CO₂ sensitivity value was overestimated and the subject was never revisited, as it should have been, the politicians have reached a solution to a nonexistent problem; but worse their solutions could actually be dangerous to humanity and the planet.


The following logistic equation is commonly found in natural biological process and economic analysis and I believe that it also applies to what we call the Greenhouse effect. My thought is to show that the base is H₂O and then add to that base a series of curves based on the CO₂ level so there would be a high and low range for temperatures for every level of H₂O. The reason I was looking for this kind of function was that if a log function is used for the sensitivity values of CO₂ then it is not asymptotic on both ends above zero. This creates problems at the low end which could not exist in the real world; therefore, it cannot be the right curve. The logistic function solves that issue.

The Following Chart is a rough approximation of my thoughts at this time for CO₂. It is based on the following Logistic function

$$Y = C / (1 + Ae^{-Bx})$$

C is the upper limit 2.7
A is the number of "doubles" to reach C 110
B is a number that controls the slope .014
X CO² ppm starting value 0

Chart 10, CO₂ Sensitivity Value Model, Logistics Function




The Following Chart is a rough approximation of my thoughts at this time for H₂O. It is based on the following Logistic function. The same logic about using a logistic function verses a log function also applies to water.

$$Y = C / (1 + Ae^{-Bx})$$

C is the upper limit 42
A is the number of "doubles" to reach C 80
B is a number that controls the slope .0025
X H_2O ppm starting value 0

Chart 11, H₂O Sensitivity Value Model, Logistics Function



The following chart **Figure 8** is from NASA and uses anomalies from the base of 14.0° Celsius. This method makes it difficult to make comparisons because the14.0° base value has no meaning and the chart itself is suspect. So we will now have to look at how the temperature is measured so we can understand the methods because the problem, intentional or not, goes back to physics and how we show information. It's critical that when we talk to nonscientists that information is properly displayed. And nowhere is this more important than when we are discussing global temperature in relationship to anthropogenic climate change.

Figure 8 is also misleading because of the methods used to make this Figure have been modified to make the look, fit their theories. The previously shown **Figure 4** shows a more reasonable chart of geological temperatures and CO₂ values than the misleading one shown in **Figure 8** especially over the past ten thousand years. The red circle shows a very missing temperature plot; look on page 20 at **Figure 9** and you can see the difference.




Figure 8, NASA Official Temperatures


When we talk about climate (long term changes; meaning centuries) or weather (short term changes; decades) local temperatures are going be in degrees Celsius (C) in the EU and science, or degrees Fahrenheit (F) in America. The base temperature for the earth that NASA established is 14.0° C or 57.2° F; but these are both relative measures and **do not tell us how much heat** (thermal energy) is there. To know that we must use Kelvin (K) or Rankin (R) and that would be 287.15° K and 516.87° R all four of those numbers 14.0° C, 287.15° K 57.2° F and 516.87° R, are exactly the same temperature, just using a different base. But if the current temperature went from 14.0° C, to 14.86° C that is a 6.14% increase in C, an increase of 2.71% in F and an increase of .30% in K and R; so which one is real? The answer is .30% because Kelvin and Rankin are the only ones that measure the total increase in energy! **Table One** shows these relationships that we just discussed.

Table One		Change in T	hermal Energ	I y
	Celsius	Kelvin	Fahrenheit	Rankin
Base, 1950 to 1980	14.00	287.15	57.20	516.87
2017	14.86	288.01	58.75	518.42
Percent Increase	106.14%	100.30%	102.71%	100.30%

The next step is to plot Carbon Diode (CO₂) from NOAA-ESRL and the estimated global temperature as published by NASS-GISS each month. As can be seen in **Table One** on the next page It doesn't

really matter whether we would use Kelvin and Rankin since the increase in thermal energy is exactly the same either way; but we'll use Kelvin as that is the accepted norm in the scientific community for determining the amount thermal energy in any object especially when looking at changes in temperature or measuring the thermal energy in any object. There are other less known temperature scales that have specific purposes, but they don't really apply here in this subject.

The important thing is how much has the global temperature actually gone up since we started to measure CO_2 in the atmosphere? To show this graphically **Chart 8** was constructed by plotting CO_2 as a percent increase from when it was first measured in 1958, the Black plot, with scale on the left shows CO_2 going up about 33.0% from 1958 to October of 2018. That is a very large change as anyone would have to agree. Now how about temperature, well when we look at the percentage change in temperature from 1958, using Kelvin (which does measure the change in heat), we find that the changes in global temperature (heat) are almost un-measurable. The scale on the right side had to be expanded 10 times (the range is 50 % on the left and 5% on the right) to be able to see the plot in the same chart in any detail. The red plot, starting in 1958, shows that the thermal energy in the earth's atmosphere increased by .3%; while CO_2 has increased by 33.0% which is over 100 times that of the increase in temperature. So is there really a meaningful link between them that would give as a major problem? The numbers tell us no, there isn't.



The next chart is Chart 8a which is the same as Chart 8 except for the scales which are the same for both CO2 and Temperature. As you see the increase in energy, heat, is not visually observably in this

chart hence the need for the previous chart 8 to show the minuscule increase in thermal energy shown by NASA in relationship to the change in CO₂. Based to these trends, determined by excel not me, in 2028 CO₂ will be 428 ppm and temperatures will be 15.0° Celsius and in 2038 CO₂ will be 458 ppm and temperatures will be 15.6° Celsius. This is what the data shows no matter what the reasons are, so I have no idea how the IPCC gets to predict that the world will end in ten or twenty years.





Chart 8 and Chart 8a are both based on the following two data series. First NASA-GISS estimates of a global temperature shown as an anomaly (converted to degrees Celsius) as shown in their table Land Ocean Temperature Index (LOTI) and as shown next in Chart 1 as the red plot labeled NASA the scale for the temperatures is on the left. The NASA LOTI temperatures are shown as a 12 month moving average because of the very large monthly variations. Second NOAA-ESRL CO₂ values in Parts per Million (PPM) which are shown in Chart 1 as a black plot labeled NOAA the scale for CO₂ is shown on the right no change is required to the NOAA data set as it is ready to use as is. The NASA data which is a mathematical construct shows there are swings in temperatures which are not believable and on the global bases that implies very large changes in the heat energy on the plant with would appear to be unrealistic. Planetary changes can just not realistically be that great on a month to month basis.

NASA published data is shown as an anomaly, but what is a temperature anomaly? An anomaly is a deviation from some fixed base value. There were two problems with the system that NASA picked which was there is no "actual" global temperature and climate is a variable so there cannot be a real base to measure from and certainly not 1950 to 1980. NASA known for its science and engineering expertise back in the day thought they could get around these issues and created a system to do so. First, they developed a computer software system they called homogenization which took all the readings from all over the planet, and then made adjustments to them in the software and then came up with the **estimated** global temperature. Second, they picked the period 1950 to 1980 (30 years) and averaged the values found in that period and came up with 14.00 degrees Celsius and made that their base. Lastly, they took the calculated monthly temperature and subtracted the base from it which gave them the anomaly after multiplying the result by 100. In **Chart 1**, we show the actual temperature not the anomaly by reversing the process. We'll talk more about this later.

Chart 1, NASA and NOAA actual numbers



The problem is that both are arbitrary. Why pick 1950 to 1980, the area between the green lines, as the base period? Is there something special about that time frame? And as to a global temperature there is no such thing for many reasons like the earth faces the sun so one side is cool and onside it warm. Higher latitudes are cooler than the equator and higher elevations are cooler than lower. And finally there are many areas where there are no measurements taken. Therefore, there is no one temperature only an artificial artifact solely dependent on the number of data points and soundness of the software used to create that one temperature! **Chart 1** above accurately show only show NASA

and NOAA data as published with no manipulation other than using a 12 month moving average for the NASA data.

As previously discussed in this paper on page 19 the current base of 14.0° Celsius was an ad hock selection of 360 values from 1950 to 1980. Using the base NASA shows global temperatures moving up slightly but the same thing could have been shown going back to any block of time. For example, the little ice age which reached its lowest temperature about between 1600 and 1650 where there are pictures of ice skating on the Thames River, in London. This shows that temperatures have been on the upswing for over 400 years now.

The little ice age is also shown in **Figure 9** which is a chart that was developed from ice core samples taken from Greenland, there have been several significant swings in temperatures over the past 10,000 years. The point being made here is what period or base should we measure the estimated temperature from as the current NASA section of 14.0[°] Celsius is just a meaningless arbitrary reference point.





Figure 9, Temperatures from Greenland ice cores

What is shown next are two charts **Charts15** and **Chart 16** that were developed to show two recommended new bases for showing the global temperature estimate as published by NASA. The first Chart, **Chart 15** uses the conventional black body temperature of the planet as the base because it is a real number that is fixed on the orbital parameters and the suns output at -18.75° Celsius (254.39° Kelvin) instead of the arbitrary 14.0° Celsius that NASA concocted. Using this number drastically simplifies the homogenization process. We don't need to discuss the details of why in this

paper. The solid blue line at the bottom is the zero point on the scale on the left side. I've also added the geological high (red), mean (green) and low (blue) for reference. This chart shows tow things number one the magnitude of the real greenhouse effect approximately 33° Celsius. And using that reference puts the current temperature value in prospective, which are that we are way to the low side of geological temperature not high side as we are led to believe.


Chart 15, NASA Anomalies using the Planets Black Body Temperature of 254.36 degrees Kelvin as the base.



The following **Chart 16** shows the same exact temperatures as **Chart 15** or **for that matter Chart 1** from the new prosed base of 17.0° Celsius which is the estimated mean temperature of the planet geologically as shown in **Figure 4**. Although I would prefer the black body temperature to be used as the base, that's probably too technical so the next best would be 17.0° Celsius the mean global temperature. And if that 17.0° isn't a good number then the science community should study it and find out what it is, without political interference.

The advantage with this kind of chart is that it shows that the current world temperatures are historically very low as they basically run from -3.0° C to -2.0° C from the global mean. Keep in mind that there is now a panic that if the published global temperatures went up 2.0° Celsius from the base of 14.0° Celsius that the planet would melt and all life would be gone. Well I'm not using new math here so 14.0° C plus 2.0° C only adds up to 16.0° C which is not even to the green line on the chart, the mean average of the planet historic temperature.

Chart 16, NASA Anomalies using the Geological Mean temperature of the Planet 17.0 degrees Celsius as the base.



What is shown here in **Chart 16** is the best base to use to measure from whether we use temperature of anomalies as used by NASA in their homogenization process and therefore the IPCC. The scientific and engineering reason is that by using the period from 1950 to 1980 the base falls into the period under evaluation and as adjustments are made every month from 1880 to the current month the base period cannot be allowed to change. So how can the base fall inside the range being measured when the entire range is recalculated every month that is using circular logic. I know I would not want to try and program that into the homogenization process.

What follows next is a page from a scientific paper written by W. A. van Wijngaarden and W. Happer published on June 8, 2020 titled *Dependence of Earth's Thermal Radiation on Five Most Abundant Greenhouse Gases*. It's a 38 page work with significant ramifications to the validity of the IPCC climate change narrative. The bottom line to this scientific study is that there is NO DANGER to additional CO₂ in the earth's atmosphere. Any warming that might be caused by CO₂ has for the most case already been accounted for.

Page 13 from that report, on the next page, clearly shows that the sun's radiation absorption bands for CO_2 are now saturated and there will be no additional effect. The green line is no CO_2 the back line is the current level of CO_2 and the red line is double the current level of CO_2 . As can be clearly seen the black and red lines are virtually identical.




Figure 4: Effects of changing concentrations of carbon dioxide, CO_2 on the filtered spectral flux $\langle \tilde{Z}^{\{i\}}(\nu, z_{\rm mp}, f) \rangle$ of (44) at the mesopause altitude, $z_{\rm mp} = 86$ km. The width of the filter (43) was $\Delta \nu = 3$ cm⁻¹. The smooth blue line is the spectral flux, $\tilde{Z} = \pi \tilde{B}(\nu, T_0)$ from a surface at the temperature $T_0 = 288.7$ K for a transparent atmosphere with no greenhouse gases. The green line is $\langle \tilde{Z}^{\{i\}}(\nu, z_{\rm mp}, 0) \rangle$ with the CO_2 removed but with all the other greenhouse gases at their standard concentrations. The black line is $\langle \tilde{Z}^{\{i\}}(\nu, z_{\rm mp}, 1) \rangle$ with all greenhouse gases at their standard concentrations. The red line is $\langle \tilde{Z}^{\{i\}}(\nu, z_{\rm mp}, 2) \rangle$ for twice the standard concentration of CO_2 but with all the other greenhouse gases at their standard concentrations. Doubling the standard concentration of CO_2 (from 400 to 800 ppm) would cause a forcing increase (the area between the black and red lines) of $\Delta F^{\{i\}} = 3.0$ W m⁻², as shown in Table 2.

The effects on radiative transfer of changing the column density of the *i*th greenhouse gas to some multiple f of the standard value, $\hat{N}_{\rm sd}^{\{i\}}$, can be displayed with filtered spectral fluxes


$$\langle \tilde{Z}^{\{i\}}(\nu, z, f) \rangle = \langle \tilde{Z}(\nu, z, \hat{N}_{\text{sd}}^{\{1\}}, \dots, \hat{N}_{\text{sd}}^{\{i-1\}}, f \hat{N}_{\text{sd}}^{\{i\}}, \hat{N}_{\text{sd}}^{\{i+1\}}, \dots, \hat{N}_{\text{sd}}^{\{n\}}) \rangle. \tag{44}$$

Figs. 4 and 5 show how varying the concentrations of CO_2 and CH_4 affect the filtered spectral fluxes at the mesopause altitude, $z_{\rm mp}=86$ km. Expanded views of the differences between the flux for standard and doubled concentrations of greenhouse gases are shown in Fig. 6, where we display

$$\langle \Delta \tilde{F}^{\{i\}}(z_{\rm mp}, 2) \rangle = \langle \tilde{Z}^{\{i\}}(\nu, z_{\rm mp}, 1) \rangle - \langle \tilde{Z}^{\{i\}}(\nu, z_{\rm mp}, 2) \rangle. \tag{45}$$

Part Three, the Probable Range of Temperatures on the Planet

The next Chart, **Chart 12** is derived from **Chart 10** and **Chart 11** and is created using the following logic. The first curve is the dark blue line at the bottom of the S shaped curves which run across **Chart 12** from left to right represents the equation for H_20 previously shown except we start at -18.89 C which is the accepted blackbody value of the Earth. The curve shown here is therefore the greenhouse effect of H_20 with no CO_2 present; we are ignoring other gases as their contribution is minimal at present concentrations. H_20 is on average 2,500 ppm and that is where the purple vertical line is placed; and that vertical line intersects the dark blue line at about 12 degrees C which just happens to be very close to the lowest estimate for the planets geological temperature as shown by the graphic on page 5 and shown here as a Cyan line. The Red line is the Global max temperature 22° C, and the Green line is the current global temperature of about 14.9° C.



Next, we add to the base H_2O line, lines for CO_2 at various levels in the atmosphere. The core assumption is that as CO_2 level go up the global temperature will follow by transferring energy to the water the additional heat may increase the level of H_2O in the atmosphere but there is no evidence

that this varies much at a global level; probably less that a 100 ppm. But both H₂O and CO₂ have saturation limits based on the parameters set in the individual curves and so no runaway effect is possible. This conclusion is supported by geological records that indicate the global temperature has ranged about 10 or 12 degrees Celsius and CO₂ has ranged about 200 to 6,000 ppm.

In the original version of this paper there were other plots shown here for different levels o co2. There was a blue line is for 300 ppm CO₂ a brown line for 400 ppm CO₂ a dashed gray line for 500 ppm CO₂ and a solid black line for 1,000 ppm of CO₂ which was the saturation point above which there is no longer any meaningful greenhouse effect. The only plot shown in Chart 12 is he brown line for 420 ppm CO₂ which intersects the horizontal green line at 15 degrees C and which is the approximate current global temperature. The reason for eliminating all the other plots was the very important paper published by Wijngaarden and Happer in 2020 (one chart is shown on page 23) which proved there was no meaningful additional captured heat past 400 ppm.

Now for a summary we have an exploded view of **Chart 12** labeled as **Chart 13** where we zoom in to get more detail of the temperature ranges of the planet based on realistic numbers and equations. The chart is based on the black body temperature of the planet, realistic logistics equations for H₂O and CO₂ and lastly geologic temperature estimates for lows, highs and a mean. None of these can be shown to be false, although I would be the first to agree that these numbers could be adjusted some by serious scientific work.

- We know that the Black Body temperature of the Earth is 254.39^o Kelvin on average but there is a variance of about 2.15^o Celsius plus or minus.
- We know that the amount of H₂O (water) in the earth's atmosphere is by far the most significant greenhouse agent and is the primary determinant of the temperature of the planet at about 85.0% of the total greenhouse effect.
- We know that the amount of CO₂ (Carbon Dioxide) in the earth's atmosphere is also a greenhouse agent but it is only a secondary determinant of the temperature of the planet at maybe 16.0% of the total greenhouse effect.
- We have developed an equation for the sensitivity values of H₂O and CO² using a logistics function rather than a log function since a logistic function more accurately represents the actual process of this kind of item as there are limits to the values both negative and positive on the planet temperature.
- We also know the probable high, low and mean geological temperatures of the planet for the past 600 million years.
- We know that the actual temperature of the planet is very stable despite all the major events that have impacted the planet which indicates that the positive and negative feedbacks are in balance.

Based on those above facts and acquired knowledge we can create a representation of all the possible stable temperatures for different amounts of H₂O and CO₂ in the planet's atmosphere and that is now shown graphically in **Chart 13**. The box represented between the vertical yellow lines and the red and blue horizontal lines contain all reasonable possible temperatures for the planet based on what we have developed in the analysis.

Chart 13, Global Temperature H₂O + CO₂ Logistics Function



The amount of water in the earth's atmosphere probably falls between 2,250 ppm (.225%), 3,100 ppm (.31%) which are represented by the two Blue dashed vertical lines, and these values are probably directly related to the planets temperature. The estimated minimum and maximum global temperatures appear to fall between 12° and 22° Celsius which falls between the Cyan and Red horizontal lines. The last thing we know is that the current temperature is about 14.9° Celsius (green horizontal line) and the water in the atmosphere is .25% which is shown as purple vertical line. The yellow line is 17° Celsius representing the mean temperature of the planet. Lastly using the logistics functions that were developed we have these curves.

- First the dark blue curve starting at the bottom left and running to the upper right which represents the planet's atmospheric temperature with no CO₂ in it.
- Second, we have the brown curve starting at the bottom left and running to the upper right which represents the planet's atmospheric temperature with CO₂ at the 420 ppm level, which is where it is today.

- In summary, the vertical purple line and the green horizontal line intersect the brown curve such that the current planet's climate/weather conditions are satisfied. The H₂O is at .25%, the CO₂ is at 418.9 ppm and the current temperature is 14.9 Celsius. The green circle.
- The 14.9° Celsius Temperature is suspect because of the homogenization process that NASA uses but that is a separate subject only briefly discussed here on Pages 9 and 10. But with no solution to the problem presented here, we really don't know what the planets temperature is with any specificity.

Now it is given that these numbers are estimates and so may very some, but we also know that the earth's temperature is inherently very stable so they cannot very much from what is shown here in **Chart 13**. The key factor is that the planets temperature is a function of the amount of water in the atmosphere and that the CO₂ levels contribute to that temperature. However since there is a saturation point to the Caron Dioxide / Water dynamic governed by the method that energy is transfer between the two it requires a higher percentage of water in the atmosphere to be able to absorb the additional carbon captured energy so the two cannot be looked at independently.


Therefore, it is my suspicion that NASA is either measuring weather or they do not understand that there are cycles to climate. This fundamental error results in an improper set of assumptions such as ignoring the variability of the Black Body temperature of the planet and assuming that CO_2 either is the primary driver of global temperatures or that it has a greater influence then it does since it is the water that actually holds the heat not the Carbon Dioxide. The core problem is the sensitivity value that the IPCC assigned to CO_2 in 1979. That value is the subject for a different paper but in general peer-reviewed papers since then have significantly reduced that value ever since. That alone destroys every IPCC global climate model.

But now that politics has gotten involved there is no hope of changing the direction as government propaganda and misinformation now drives the narrative in the elementary schools, the high schools the and colleges. It will take multiple generations to undo the damage that was done which takes it into the next century.


Starting on page 30 there are two 4 page printouts of NASA table LOTI. The first is from April 2008 which is the oldest one I saved and the second is Current date of the paper. So let's look at the numbers that NASA publishes. First on the April 2008 printout look at the anomalies for the entire year 2007 which average .539 degrees Celsius higher than the base. Then look at the same values on the August 2022 printout and we find the average for 2007 is .675 degrees Celsius.136 degrees Celsius higher than 2007 so in 12 years and 1 month the past got 25.2% warmer --- now how did that happen? The answer is simple the process that NASA uses to calculate anomalies are not stable and therefore all the numbers are a **variable**. If the past, can be 25.2% warmer after 15 years that makes the present 25.2% colder so how do we even know what the global temperature is? The answer lies in the way that NASA calculates global temperatures which they call homogenization. The bottom line is that every time they run the program they get different results. Some of these "adjustments" seem to give the impression that there is a desired result that isn't there in the raw data but by tweaking the program the raw data can be made to look like what someone wants it's to look.

So, now let's look at two more charts the first, Chart 1a, will be a plot of 12 years of temperatures (from LOTI Table) out of the past 20 years (to simplify the chart). The plot shows that, for example, for the anomalies of 1999 from the year 2012 to the current full year 2021 (Red plot with a star). In general, all the years in the NASA process almost never gives the same value in succeeding years that plot should just be a straight line. Some of these changes are large and others aren't. Some move up and some move down. Why the large jump in temperature in 2015/2016, maybe the Paris Climate change Meeting?

Now look at the plots for 2015, 2016, 2017, 2018, 2019, 2020 and 2021, the top seven plots. That doesn't seem to make sense since there was not a corresponding large jump in CO₂ so why the dramatic change?



Now let's look at another chart, Chart 1b, this one is from NOAA showing CO₂ levels in PPM. This chart was developed the same way that the previous one was and we see that the values for CO₂ do not change after they are calculated. This is the way a chart of this type should look and so there is no reason to ever show this chart, Chart 1b, of CO₂ the one on page 19 is sufficient.



So looking back to the weeks after the November 2018 election we have Alexandria Ocasio-Cortez (AOC) telling us that we only have 12 more years to live if we didn't give up all carbon based fuels that we use "immediately" so; we now have a hard date to measure from and we also know that the point of no return is an increase of 2.0° Celsius from the base of 14.0° Celsius or 16.0° Celsius. Now 12 Years from November 2018 would be November 2030 but we'll be generous and say we have until January 2031. The present global temperature is .93° Celsius above the base of 14.0° Celsius or 14.93° Celsius. This only gives us 1.07° Celsius of increase left as we have already used up 46.5% of the available buffer before we die.

Do we really believe any of this, I don't! And apparently, neither do the insurance companies or any of the communities on the shorelines around the entire planet believe this. And also I believe that the NASA numbers are manipulated to give results that the politicians want --- this entire climate change scare is all made up!

The following from Sir Karl Pooper should be understood by everyone in science as gospel.

<u>Sir Karl Raimund Popper</u> (28 July 1902 – 17 September 1994) was an Austrian and British philosopher and a professor at the London School of Economics. He is considered one of the most influential philosophers for science of the 20th century, and he wrote extensively on social and political philosophy. The following quotes of his apply to this subject.

If we are uncritical, we shall always find what we want: we shall look for, and find, confirmations, and we shall look away from, and not see, whatever might be dangerous to our pet theories.

Whenever a theory appears to you as the only possible one, take this as a sign that you have neither understood the theory nor the problem which it was intended to solve.

... (S)cience is one of the very few human activities — perhaps the only one — in which errors are systematically criticized and fairly often, in time, corrected.

David J. Pristash, Independent Researcher BBA, EMBA, Graduate GE management program, Captain US ARMY 18A (WIA Retired), Eight issued patents' Member Beta Gamma Sigma Brecksville Ohio 44141 Email David.Pristash@gmail.com Face Book www.facebook.com/david.pristash Blog www.centinel2012.com Cell 216 272 4583

GLOBAL Land-Ocean Temperature Index in 0.01 degrees Celsius base period: 1951-1980

sources: GHCN-v3 1880-11/2012 + SST: 1880-11/1981 + HadISST1 12/1981-11/2012 Reynolds v2 using elimination of outliers and homogeneity adjustment

Notes: 1950 DJF = Dec 1949 - Feb 1950 ; ***** = missing

April 2008

base period: 1951-1980 GLOBAL Land-Ocean Temperature Index in .01 C

sources: GHCN 1880-04/2008 + SST: 1880-11/1981 HadISST1 12/1981-04/2008 Reynolds v2

using elimination of outliers and homogeneity adjustment Notes: 1950 DJF = Dec 1949 - Feb 1950; **** = missing

	Year	∞	88	∞	88	88	88	88	88	1888	88	89	89	89	89	89	89	89	89	89	89	90		69	90	90	90	90	90	90	90	90	90	1910	91
	SON	-26	-31	-30	-29	-33	-26	-26	-36	-12	-29	-39	-28	-33	-21	-35	-14	-10	-17	-31	-1	8		SCS	-25	-34	-39	-27	-18	-25	-33	-32	-23	-35	-21
	JJA	-26	-18	-24	-13	-30	-31	-16	-21	-24	-17	-33	-23	-28	-21	-29	-22	-13	8	-18			ļ	OOA	-17	-25	-37	-34	-21	-15	-38	-27	-31	-28	-30
	MAM	-26	-4	-20	-19	-34	-30	-21	-34	-35	-4	-36	-23	-36	-28	-31	-29	-27	8 1	-36	-22	9-	,,,,,	MAM	-4	-28	-29	-37	-26	-16	-38	-41	-43	-33	-45
	DJF	****		6-	4	-20	-41	-31	4	-41	6	3	-44	N	9	3	-45	-19	-16	٦	-28	2	- 1		-	-		4	3	N				-33	
ean	D-N	*	-19	-21	-26	-29	-32	-24	-34	-28	-15	-36	-30	-29	-33	-33	-28	-17	-12	-24	-17	-11		Z C	-14	-25	-31	-36	-25	-21	-36	-34	-35	-32	-37
AnnMean	J-D	-25*	2	2	N	3	3	2	3	-26	\leftarrow	3	2	3	3	3	N	\neg	-1	N	-17	-10		1	\vdash	S	-31	3	N	S	3	-34	-35	-33	-34
	Dec	-22	-25	-48	-23	-32	-13	-28	-38	-21	-29	-35	-13	-47	-34	-30	-20	-13	-12	-26	-28	-10	ſ	Dec	-28	-46	-42	-20	-20	-12	-44	-40	-40	-56	-19
	Nov	-30	-33	-30	-30	-34	-27	-25	-39	-3	-35	-52	-40	-42	-21	-39	-12	-17	-19	-39	13	-16	;	NON	-24	-42	-31	-11	8	-35	-43	-41	-20	-41	-16
	Oct	-27	-29	-36	-29	-33	-26	-34	-43	-12	-29	-26	-25	-33	-17	-29	-17	-2	-18	-32	-5	-2		OCT	-24	-32	-42	-30	-28	-15	-26	-33	-23	-32	-21
	Sep	-20	-29	-24	-29	-33	-27	-20	-26	-21	-23	-38	-19	-23	-24	-37	-13	-13	-15	-21	-12	-7	Ţ	Sep	-26	-29	-45	-41	-19	-25	-29	-23	-25	-30	-25
	Aug	-19	-13	-16	-18	-26	-27	-18	-29	-25	-22	-33	-22	-29	-25	-25	-21	-14	8	-16	-11	-11	,	And	-19	-28	-39	-31	-18	-10	-37	-33	-24	-26	-29
	Jul	-21	-10	-27	-10	-27	-24	-2	9	-22	-17	-29	-26	-33	-11	-20	-21	-10	-2	-17	-14	-11	r	Jul	-18	-18	-28	-34	-19	-21	-37	-22	-33	-21	-25
	Jun	-40	-33	-30	6-1	-36	-41	-29	-25	-25	-11	-36	-22	-21	-26	-41	-24	-14	-15	-20	-23	8-	ŀ	Jun	-15	-29	-43	-37	-25	-14	-41	-26	-37	-36	-36
	May	-30	-4	2	2	3	3	\vdash	-25	2	-5	4	N	3	3	3	-30	-15	H	-34	-19	91		ď	-	3	-34	3	2	2	4	3	3	-34	4
	Apr	-25	9	2	Н	3	3	\vdash	-39	3	-2	3	-31	4	3	3	2	-38	15	-25	-20	-111		Apr										-30	4
	Mar	-23	-2	6-1	\vdash	3	-19	3	3	-46	-4	-31	\vdash	3	\vdash	N	-30	2	-18	-48	-27	-2		Mar	_	-24	-14	-30	H	-21	-25	-48	-45	-34	-48
	Feb	-20	-22	9	3	-	3	3	5	4	œ	-41	5	8	-54	-34	-53	-16	-14	-23	-37	7	32		-		2	4	5	3	-45	2	3	-30	-46
	Jan	-23	-28	-7	-41	2	9	4	9	3	-	-45	4	3	-87	4	5	-20	2	9	2	-32		Jan	2	-16	-24	-51	-30	-30	-45	4	-58	-28	-52
	ea		88	88	88	88	88	88	88	∞	88	89	89	89	89	89	89	89	89	89	89	90		e B	90	90	90	90	90	90	90	90	90	1910	91

7 8 4 5 9 7 8 6 0	(m 4 5 9 7 8 6 0	7112 M 4 5 9 7 8 6 0
1913 1914 1916 1917 1919 1919	9000000 900000000000000000000000000000	000000000000000000000000000000000000000	11933 11933 11933 11933 11933 11933	Year 1941 1942 1944 1946 1947 1950
-45 -27 -11 -12 -30 -30 -24		1 1 1 0 0 1 1 7	-18 -11 -11 18 17 -17	SON 116 222 222 16 16 19 19
138 17 17 17 17 17 17 17 18	JJA -111 -21 -25 -19	-111 -25 -25 -12	-10 -22 -11 -11 -3 -3 -3	JJA 8 8 2 9 1 1 6 7 7 7 7 1 1 6 1 6 1 7 7 7 7 7 7 7
26 36 30 30 47 15	MAM -114 -23 -20 -20 -19	11111111	116 116 118 118 118	MAM 10 3 7 18 10 0 6 -2 -7
1 1 3 6 1 1 1 4 1 1 1 3 6 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		1001	110 110 110 110 110 8	DJF 155 123 33 30 144 -10 -10
1 3 3 6 8 8 4 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9		11 11 2 11 1	115 129 112	D-N 10 20 22 22 22 -23 -13
134 131 151 130 130 132 120 190 190 190	O 8 4 L L 9 L	111 22 25 113 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	73088765	J-D I 111 110 211 21
-35 -15 -15 -20 -66 -70 -32	2 H M I H H G	12 4 1		Dec 22 22 6 13 13 13 10 10 10
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Nov -177 -13 -13		-25 -22 -22 -20 -20 -20	No 4 12 1 19 19 19 19 19 19 19 19 19 19 19 19 1
-53 -32 -32 -23 -17 -17	Oct - 25		12 0 2 16 16 22 22	0ct 21 23 24 25 25 25 13 13 13 13
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Sep 		113 113 114 114	Sep - 12 - 12 - 2 - 2 - 2 - 1 - 1
- 53 - 13 - 13 - 15 - 15	Aug -24 -25 -25 -12		-10 -11 -2 -11 -1	Aug 2 -1 6 17 16 -18 -3 -4
-39 -17 -17 -27 -18 -29		-11 -24 -6 12	-4 -0 11 11 8 6	Jul 13 13 16 16 16 19 19 19
-23 -20 -20 -34 -34 -15	Jun -24 -22 -19	-15 -27 -29 -14	-16 -13 -13 -2 -2	Jun 98 8 8 8 8 12 12 12 13 -7 -7 -5 -55
	May - 188 - 29 - 20 - 23	1 1 3 3 7 1	-17 -21 -9 -9 -0	May 10 10 17 17 -6 -6 -7
-20 -20 -27 -26 -40 -10			122 122 123 123 123 123	Apr 151 19 19 19 10 10 10 10
-35 -35 -133 -133 -27 -27	Mar 113	- 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	-116 -123 -134 -133	M ar 122
-116 -111 -111 -16 -53 -19		-4 -42 -13 -18	-17 26 -26 -26 11 11	Feb 22 22 22 26 13 13 13 9 9 15 15 15
	Jan -31 -24 -28	- 19 - 19 - 14 - 14	-20 -14 -25 -19 -16	Jan 11 22 -12 37 16 14 -24
1912 1913 1914 1915 1916 1917 1918 1919	90000000000000000000000000000000000000	933222	1933 1934 1935 1936 1937 1939 1940	Year 1941 1942 1943 1944 1945 1947 1949

1951 1952 1953 1954 1955 1955 1956 1959	0000000	1968 1968 1970 1971 1972 1973	0000 000 000 000 000	Year 1981 1982 1988 1984 1986 1987 1988
117 - 13 - 0	SON 7 7 16 -29 -10 -10		-111 -17 18 18 15	SON 12 23 23 6 3 19 22
27	JJA 6 0 16 -10 -13		1 1 2 1 1	50 29 29 21 21 21 21 21 21 21 21 21 21 21 21 21
- 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1	MAM 14 5 - 29 - 12	0 4 0 6 4 1 2 4 1 6	-24 -21 -24 -24 -24	MAM 28 28 31 17 18 18 18 18 15 15
27 - 15 - 15 - 27 - 27 - 27 - 27 - 27 - 27 - 27 - 2	71 11	-112 -122 -111 -119 -16	14 10	DUF 31 12 12 38 38 13 72 22 23 43
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	D-N 10 3 8 -18 -13	10 6 4 4 11 11 11 11 11 11 11 11 11 11 11 11	7 1 1	D-N 25 28 28 11 13 13 19
-4 -111 -100 -177 -177 -17	J-D 8 4 4 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	0 0 4 8 8 8 1 8 1 8 1	1	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
118 117 117 17 17 17 17		112 113 113 113 113 113 113 113 113 113	7 3	Dec 29 34 11 11 7 7 43 118
1133 122 122 123 124 126 127 127 127 127 127 127 127 127 127 127		2-11001-	7 7 7 7 7	NO N 1 8 4 4 8 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	oct 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	113 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1	1211	0ct 102 102 50 209 209
0 8 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Sep 8 2 2 -37 -16	0 0 4 4 4 1 1 2 1 0 0 1	1 - 1	Sep 12 13 13 13 13 13 32 32
110 100 111 112 122 126	Aug 3 - 5 - 24 - 6	-5 -12 -12 -20 -11	-21 -19 -21 -21 13	Aug 35 30 13 11 11 15 28
11	Jul 1 -3 14 -5 -21	6 7 1 1 1 1 1 1 1 1	13 20 20 7 18	Jul 31 13 13 14 -3 6 26
1 1 1 1 2 4 4 1 1 1 1 2 4 4 1 1 1 2 2 1 1 1 2 2 1 1 2 2 1 1 2 2 1 1 2 2 1	Jun 13 7 6 -2 -12	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	- 14 - 23 - 8 - 8 - 10	Jun 20 00 18 12 12 6 39
11 118 122 122 10 10	May 22 - 10 - 28 - 28 - 6	111 - 133 - 111 - 125 - 127 - 1	18 -29 -29 -1 -6 -27	May 115 114 30 30 30 12 12 37
- 10 - 10 - 110 - 21 - 21 - 10 - 113	Apr 11 11 12 -7 -19	21 - 20 - 21 - 21 - 21 - 21 - 21 - 21 -	-16 19 11 23	Apr 24 24 24 25 26 2 2 2 2 35 35 11 35 15
-177 -18 -29 -22 -32 -32 -32	Mar 10 13 -13 -10	1 2 1 2 4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	11 -27 15 12 9	Mar 433 37 21 12 25 16 47
139 144 124 123 123 18		123 123 120 120 120 120 120	7 1 7 7	Feb 37 37 37 98 38 83 83 85 85 85 85 85 85 85 85 85 85 85 85 85
-32 -14 -18 -19 -15 -15 -111	Jan 5 1 1 -10	100	1 10 3 6 21	Jan 44 47 123 15 20 51 3
1951 1952 1953 1955 1955 1956 1959 1959	90000000000000000000000000000000000000	1967 1967 1970 1971 1972 1973	7 C C C C C C C C C C C C C C C C C C C	Year 1981 1982 1984 1985 1986 1987 1988

1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	Year	2001	2002	2003	2004	2005	2006	2007	2008	Year
32	26	8	9	34	35	26	49	42	30	25	SON	53	50	59	26	89	59	52	***	SON
35	46	9	ω	25	41	33	38	67	31	35	JJA	47	49	20	32	57	52	54	* * * *	JJA
51	34	23	21	21	30	26	37	58	25	42			99						*	MAM
29	39	33	23	80	46	33	29	61	20	34			64						*	DJF
37	36	13	14	22	38	29	38	57	34	34	D-N	45	57	52	20	62	53	09	* * *	D-N
38	32	13	14	24	38	30	40	57	33	33			26						*	J-D
37	24	12	9	25	25	32	53	51	33	20	Dec	51	36	89	51	29	69	40	* * * *	Dec
41	19	9	m	33	37	35	99	43	32	26	Nov	19	51	49	63	64	62	49	****	Nov
40	21	-4	15	37	44	17	20	40	31	19	Oct	44	49	99	58	71	09	52	*****	Oct
17	39	-10	0	30	24	25	41	43	27	31	Sep	48	48	09	46	89	55	51	*****	Sep
27	38	7	3	18	37	44	37	63	28	38	Aug	45	45	63	43	56	58	57	*****	Aug
47	49	0	12	21	50	38	26	71	30	33	Jul	20	56	49	22	55	43	53	*****	Jul
31	49	16	10	35	34	17	20	19	36	35	Jun	47	46	39	33	59	53	53	*****	Jun
37	30	19	17	17	8	19	32	61	21	29	May	51	56	51	37	55	42	99	****	May
48	44	15	16	27	39	25	34	26	27	52	Apr	39	28	49	52	64	46	64	41*	Apr
L 9	29	35	29	20	44	33	46	99	27	46	Mar	54	84	51	58	70	55	09	09	Mar
29	45	35	28	-5	71	47	30	79	09	51	Feb	41	70	51	67	56	58	63	26	Feb
32	35	39	28	24	43	26	27	52	40	17	Jan	38	71	65	52	69	43	98	13	Jan
1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	Year	2001	2002	2003	2004	2005	2006	2007	2008	Year

Divide by 100 to get changes in degrees Centigrade. Multiply that result by $1.8\,(=9/5)$ to get changes in degrees Fahrenheit.

so add that to the temperature change if you want to use an absolute scale (this note applies to global annual means only, J-D and D-N !) Best estimate for absolute global mean for 1951-1980 is 14C = 57.2F,

	or .72F	57 92F
	or	7
	.40C	
Table Value	change:	neam [elinne
!		scale if global
ple		מותניט
Example		מלמ

abs. scale if global annual mean

GLOBAL Land-Ocean Temperature Index in 0.01 degrees Celsius base period: 1951-1980

```
sources: GHCN-v3 1880-11/2012 + SST: 1880-11/1981 HadISST1 12/1981-11/2012 Reynolds v2 using elimination of outliers and homogeneity adjustment
```

Notes: 1950 DJF = Dec 1949 - Feb 1950 ; ***** = missing

Date of this paper

base period: 1951-1980 GLOBAL Land-Ocean Temperature Index in 0.01 degrees Celsius

sources: GHCN-v4 1880-10/2022 + SST: ERSST v5 1880-10/2022 using elimination of outliers and homogeneity adjustment Notes: 1950 DJF = Dec 1949 - Feb 1950; ***** = missing

JO 415.98

Year 1880 1881 1882 1883 1884 1885 1886 1889 1890 1891 1895 1895 1896	Yea 1900 1900 1900 1900 1900 1900 1900 190
SON -18 -17 -17 -18 -27 -25 -25 -24 -19 -19 -13	SON -22 -30 -46 -37 -16 -34 -44 -29 -29 -29
7048 -158 -144 -27 -28 -27 -27 -27 -27 -21 -29 -18	-10 -15 -29 -40 -49 -25 -25 -25 -36 -36 -40 -49 -49 -49 -49 -49 -40 -40 -40 -40 -40 -40 -40 -40
MAM -111 - 6 - 15 - 37 - 23 - 27 - 23 - 27 - 23 - 27 - 27	AM A
**** -16 -16 -33 -10 -34 -34 -37 -37 -15 -15 -15	N 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
AnnMean J-D D-N -16 *** -7 -8 -10 -8 -17 -19 -28 -26 -33 -34 -31 -29 -36 -35 -17 -19 -17 -19 -22 -24 -21 -21 -22 -23 -11 -11 -11 -1	- 8 - 10 D- 115 - 115 - 115 - 115 - 115 - 127 - 227 -
0.7.9.0.0.0.0	3111111111111
2079 00 00 00 00	Dec
	No 1119
0ct 0ct 122 - 22 122 - 24 122 - 24 123 - 23 124 - 21 127 - 24 128 - 27 138 - 27 138 - 27 138 - 23 138 - 21 138 -	- 20 - 20 - 20 - 20 - 20 - 20 - 20 - 20
Sep -13 -14 -14 -14 -28 -28 -23 -23 -23 -11 -23 -25 -15 -16 -28 -15 -16 -28 -17 -17 -18 -16 -17 -17 -17 -17 -17 -17 -17 -17	Se
Aug — — — — — — — — — — — — — — — — — — —	Aug -20 -20 -45 -48 -20 -19 -34 -45 -45 -43 -33
Jul -17 -15 -30 -33 -17 -25 -10 -27 -17 -17 -17 -17 -27 -17 -27 -17 -27 -17 -17 -17 -17 -17	Jul -14 -27 -34 -26 -26 -26 -34 -34 -41 -41
Jun Jun -20 -20 -20 -20 -21 -24 -25 -25 -25 -25 -25 -25 -25 -25 -25 -25	Jun -12 -29 -41 -48 -28 -51 -38 -51 -51 -51 -51 -45 -45 -45 -45
May -9 -9 -9 -9 -9 -9 -9 -9 -9 -9 -9 -9 -9	May -114 -131 -139 -251 -251 -37 -51 -254 -51 -254 -51 -251 -251 -251 -251 -251 -251 -251
Apr -15 -15 -18 -18 -29 -27 -27 -27 -27 -27 -27 -27 -27 -27 -27	4pr -27 -27 -40 -50 -50 -50 -50 -51 -44 -442 -558 -17 -558 -17 -558 -17 -558 -17 -558 -17 -558 -17 -559 -17 -559 -17 -559 -17 -559 -17 -559 -17 -559 -17 -559 -17 -559 -17 -559 -17 -559 -17 -17 -559 -17 -17 -17 -17 -17 -17 -17 -17 -17 -17
Mar	ar. 6 6 27 22 22 22 22 22 22 22 22 22 22 22 22
Feb - 23 - 136 - 147 - 136 - 147 - 147 - 147 - 156 - 156 - 156 - 156 - 167 - 168 - 1	eb 111 111 111 111 111 111 111 111 111 1
Jan -18 -18 -28 -28 -43 -71 -34 -41 -29 -81 -29 -81 -29 -14 -29 -14 -29	an 117 117 223 34 88 84 44
880 881 382 382 383 384 884 887 887 899 90 90 91 91 90 90	H = 0 0 0 = 10 10
183 183 183 183 183 183 183 183 183 183	Year 1900 1900 1900 1900 1900 1910 1910 191

1914 1915 1916 1917 1918	Year 1922 1922 1928 1928 1929 1929 1929 1931 1931 1931 1933 934 934 944 945 945	5 5 5
-111 -33 -29	SON SON SON 112 111 111 112 113 114 114 115 116 117 117 117 117 117 117 117 117 117	1 1 1 1
-21 -18 -38 -33	222 330 330 330 330 330 330 330 330 330	
-24 -30 -57 -20	; \$\int 4\pi 0 \cap 8 \tau \cap C \cap \cap \cap \cap \cap \cap \cap \cap	1
-2 116 67 24	ZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZ	1 1 1
1 1 1 1		MH
5 – 15 4 – 12 6 – 31 6 – 47 6 – 47 7 – 27	D-1-2222 22333226333333226333333226333333322633333333	\vdash
1146		
-21 -81 -67 -29	Dec 	9 01 10
-15 -12 -46 -33 -41	1143 127 129 131 143 154 175 175 175 175 175 175 175 175	⊣ 1
-3 -24 -44 -6 -6	cct 1359335665857491177	1 1
16 20 36 22 22 117		
-15 -22 -28 -222 -331 -26 -26 -26	See 1111 See	
1 1 1 1 1 1	A L L L L L L L L L L L L L L L L L L L	5 5
25 -2 2 -11 3 -2 6 -3 6 -3 6 -3 6 -3	Jul 13 13 13 13 14 4 15 15 15 15 15 15 15 15 15 15 15 15 15	7 7
20 -2 -6 -2 -4 -4 -5 -4 -5 -3 -3 -3 -3	Jun 127 - 278 - 298 - 298 - 298 - 298 - 298 - 298 - 298 - 298 - 298 - 298 - 217 - 21	12
0 1 0 1 4 2 7		11
-29 -30 -55 -44 -12	Appr - 30 - 23 - 41 - 26 - 27 - 28 - 41 - 28 - 28 - 41 - 28 - 28 - 41 - 28 - 28 - 29 - 20 - 20 - 21 - 31 - 20 - 20	19
-23 -28 -63 -24 -21	Mar 15 - 23 - 15 - 28 - 28 - 25 - 25 - 25 - 25 - 25 - 2	11
-9 -14 -63 -24	Feb 174 - 17	1.10
5 -20 -12 -57 -47 -20	Jan F 10 Jan F 23 Jan F 24 Jan F 26 Jan F 27 Jan	7
1914 1915 1916 1917 1918 1919	ear 9921 9922 9923 9924 9925 9925 9925 9927 9927 9927 9927 9927	
	1992 1992 1993 1993 1993 1994 1944 1944 1948 1949 1949 1949 1949	7

1954	95	95	95	96	0	96	96	96	1964	96	96	96	96	96	97	97	97	97	97	97	97	97	97	97	98	T U	98	98	98	98	98	98	98	98	98	99	99	1992	99
1 1 4		\vdash		0	SON		M	16	-27	6-	_7	1	-5	10	2	9-	4	ω		-10			∞		21	SON	\vdash	15	28	14	10	10	32	29	28	38	34	$^{\circ}$	13
1 1 8			7	-2	JJA		Н	11	-10	8	0	-2	-10	\vdash		8-		12				N				JJA	3	8	25	14	12	13	33	37	28	39	46	14	20
-16	V	2	12		MAM	11	∞		-26			4										28				MAM	35	12	34	22	15	25	23	46	27	09	40	36	31
-10			2	4	DJF	\leftarrow	Π	4		\vdash	-10	\vdash	\vdash	\vdash	18	\vdash	-16	27	-14	8		17		2	39	DJE	3	21	46	21								40	
3 -11	7			I	D D-N				0 -18	1	1	1	I			ı	ı	П	I	I	-1	\vdash		П	2	0	\sim	1	\mathcal{C}	\vdash	l	\vdash	2	4	2	4	4	23	2
	\dashv		. ,	ì	J-D		()	.,	\sim	-1	Ĭ	1	ĩ	Δ,	(-)	ĩ		1(1	1	-1(18	[-		26	J-D	\sim	14	31	16	12	18	32	36	27	45	40	22	23
-18				19	Ψ	-16	8	8	-30	8	-2	-5	-14	24	-12	6-1	18		8	-17	11	ĸ	80		22	Dec	4	43	16	-4	14	13	46	29	37	40	32	21	18
125	\dashv		ω 1		Nov	Υ	9	15	-21	9-	-1	-5	-5	12	2		2	2	8	-17	9-	16	14	28	30	Nov	23	18	30	7						46		∞	4
1-2	V	4		9	Oct	0	7	15	-32	-5	-16	6	0	0	∞	-4	8	10	9-	\vdash	-24	∞	∞	26		Oct	12	13	17	14	12	15	33	38	29	44	29	9	23
110	-	-2			Sep	0	0	18	-29	-15	-3	9-	-19	8	12	9-	2	6		-2	9-	2	9	25		Sep	15	14	37	21	13	M	35	37	34	23	44	-1	12
17	1 4			2	Aug	П	-1	22	-22	-4	8	J	6-1	M	-10	1	16	2	10	-17	\vdash	18	-13	17	18	Aug	35		35									ω	11
111) (1	2		-4	Jul	1	2	9	-4	-13	ω	2	-13		П					-7			4		22	Jul	32	15	18	19						45		ω	25
-19	H —			-4	Jun	11	3	2	-4	8	П	8	6-	m	1	-17	4	19	-5	1	-12	27	-1	14	20	Jun	29		22	2	15	12	35	40	17	37	53	26	23
-20	J		4		May	12	9-	- 1	-25		$\overline{}$	\vdash	-14	18	-4	1	-3	23	-4	16	-21	33	0	M	32	Мау	25	18	34	33	14	21	25	44	17	45	34	31	28
-14 -22 -28	1	, H		-15	Apr	13	2		-32			-5	9-	17	2		0	27	-11	4		26	17	15	30	Apr	32	15	28	9	12	22	24	43	29	26	51	27	28
-15 -32 -21	1 1		- (Mar	0	\Box		-21	\Box	4	2	20	П ,	1	-18	7	29	-5	13	-21	24	19	19	29	Mar	48	M	42	26	17	30	18	51	36	0 0	35	48	36
-10 -16 -24	1 1				Feb	19	15	\vdash	-10	\vdash	1 (\mathcal{C}	-14	\dashv	N	-16	\vdash	33	\sim	∞	9-	22	10	-10	30	Feb	42	16	43	14	-4	3./	43	44	30	44	50	40	37
13	1 1			0	Jan	7	2	e - 3	0 0	ω (Ι	-19		-25		∞ (1 (-22	\mathcal{C}	-10	10	۳ ا	18	9	ω	30	Jan	23	2	53	31	22	56	32	2.7	17.	4 I	43	48	35
1954 1955 1956	95	95	200	30	σ	96	96	96	1964	900	900	96	96	9 1 0	9	200	50	97	9.7	97	97	97	97	97	∞	еа	8	86	∞	300	800	ω ω 0	ω ω ω	ω 0	ω (C	\mathcal{L}	\mathcal{L}	1992	0

1994 1995 1996 1997 1999	Yea 2000 2000 2000 2000 2000 2000 2000 20	2022 Year
30 4 4 5 8 8 8 9 8 9 8 9 8 9 8 9 8 9 8 9 8 9 8	SON 58 58 62 62 61 77 77 77 77 70 88 88 88 88 88 99 99	***
32 45 43 43 69 35	JUA 53 53 62 62 63 63 66 69 69 69 78 88 88 88 88 88 88 88 88 88 88 88 88	*
33 40 31 40 65 49	MAM	90 MAM
16 32 36 36 68 56	H 0 0 0 0 H 10 H 0 7 10 H	
30 44 44 61 40	D-N 51 64 59 66 63 66 63 66 64 74 74 74 65 65 65 65 65 65 74 74 74 74 74 74 74 74 74 74 74 74 74	* * N – Q
32 45 46 61 38 39	J-D 533 623 625 647 654 655 677 774 774 677 677 677 677 677 677 677	* 0
38 26 37 59 55 41 28	Dec 568 444 755 50 688 779 688 455 699 788 116 86 93 93 93 85 85 85 85 85 85 85 85 85 85 85 85 85	k U
44 38 64 37 30	Nov 72 72 72 73 74 73 75 81 81 88 84 66 91 88 88 82 99 99	
42 19 61 41 34 26	Oct 50 54 72 60 75 69 65 70 65 70 65 70 88 88 89 89	о то
32 34 25 52 42 38	Sep 52 63 63 62 62 71 72 77 77 77 77 77 77 85 99 99	ep
21 45 48 41 65 31	Aug Sanga Aug Aug Aug Sanga	s bn
30 37 34 66 38 39	Jul 59 61 61 26 60 60 60 73 60 60 73 82 82 82 82 94 94	
44 43 25 54 77 36	Jun 52 53 48 43 64 65 64 65 64 65 67 77 77 91 52 64 69 67 77 91 52 64 65 69 60 60 60 60 60 60 60 60 60 60 60 60 60	
28 27 34 68 36 36	May 57 64 61 37 63 47 63 47 63 75 75 80 80 95 95 82 82 82 82 82 83	>
41 33 34 64 32 57	Apr 50 58 55 61 67 47 76 76 72 72 72 79 101 110 113 Apr 1	C.
29 32 52 63 32 55	Mar 55 88 88 63 63 74 63 74 92 65 67 116 117 Mar 88	ar
79 46 41 88 64 56	6 c c c c c c c c c c c c c c c c c c c	leb N
26 24 30 58 29 25	Jan 45 77 75 75 58 74 56 102 31 64 75 76 85 117 102 81 93 116 Jan Figure 193	Jan E
1994 1995 1996 1997 1998 2000	Year 2001 2002 2003 2003 2004 2005 2004 2005 2009 2000 2010 2011 2012 2013 2015 2016 2017 2019 2020 Year 2022	ear

Divide by 100 to get changes in degrees Celsius (deg-C). Multiply that result by 1.8 (=9/5) to get changes in degrees Fahrenheit (deg-F). 40 0.40 deg-C or 0.72 deg-F Table Value : change :

Example